## Henry Ford Health System

690 Amsterdam St., Detroit MI 48202 Ph: 586-247-2992

www.KelechiOkorohaMD.com

## AMZ (ANTEROMEDIALIZATION/TIBIAL TUBERCLE OSTEOTOMY

Rehab Protocol

	WEIGHT BEARING	BRACE	ROM	EXERCISES
PHASE I 0-2 weeks	Heel touch only	On at all times during day and while sleeping*	0-90° at home	Calf pumps, quad sets SLR in brace, modalities
		Off for hygiene and exercises		
PHASE II 2-6 weeks	Heel touch only	Off at night	Maintain full extension and progress flexion	Progress non-weight bearing flexibility
		Open 0-90°		Begin floor-based core, hip and glutes work
				Advance quad sets, pat mobs, and SLR
PHASE III 6 weeks - 8 weeks	Advance 25% weekly and progress to full with normalized gait pattern	None	Full	Advance closed chain quads, progress balance, core/pelvic and stability work
				Begin stationary bike at 6 weeks
				Advance SLR, floor-based exercises, hip/core
PHASE IV 8-16 weeks	Full	None	Full	Progress flexibility and strengthening, progression of functional balance, core, glutes program
				Advance bike after 12 wks
				Outdoor cycling, elliptical, swimming after 12 wks
PHASE V 16-24 wks	Full	None	Full	Maximize single leg dynamic and static balance
				Glutes/ pelvic stability/core + closed-chain quad program and HEP independent
				Begin training sport-specific drills as tolerated after 16 weeks once cleared by MD

<sup>\*</sup>Brace may be removed for sleeping after first post-operative visit (day 7-10)



