

Complex Anterior Shoulder Stabilization Rehabilitation Protocol (revision arthroscopic repairs, MDI, open capsular shifts, latarjet)

Phase	Goals	Precautions/Restrictions	Treatment
Weeks 0-6	<ul style="list-style-type: none"> Protect surgical site and repair Decrease pain and inflammation PRICE principles Minimize muscle atrophy Maintain elbow, wrist and hand function 	<ul style="list-style-type: none"> No shoulder motion No lifting Arm to be touching abdomen when out of sling Wear sling with pillow except for hygiene and exercise performance 	<ul style="list-style-type: none"> Shoulder arm hang exercises AAROM to AROM of elbow, wrist and hand with arm in plane of body Scapular retraction isometrics with immobilizer on Core activation with immobilizer on Cryotherapy: 5-7times per day May initiate cardiovascular exercise (bike) beginning week 2 Initial visit: FOTO, QuickDASH
Weeks 6-16	<ul style="list-style-type: none"> Maintain integrity of repair Initiate PROM and slowly advance to AAROM to AROM Functional AROM of shoulder by week 16 Functional scapular mechanics by week 16 Improve motor control Improve total arm strength 	<ul style="list-style-type: none"> May discontinue pillow at week 6, and wean out of sling as able over next 1-2 weeks Do not force motion No anterior shoulder stretching or subscapularis stretching until week 8 No weight bearing through shoulder until week 12 Avoid RTC pain with strengthening 	<ul style="list-style-type: none"> Week 6: PROM-AAROM-AROM of shoulder ER/IR, flexion, & abduction <ul style="list-style-type: none"> Avoid compensation Week 8: Initiate IR/ER isometrics and scapular stabilizer strengthening Week 10: Progress strength of scapular stabilizers, RTC, forearm and core Week 12: Initiation of plyometric exercise Week 12: May begin jogging/running Week 12: FOTO, QuickDASH
Weeks 16-20) (Months 4-5)	<ul style="list-style-type: none"> Maintain integrity of repair Progress RTC exercises Progress scapular stabilizer strengthening Full AROM compared bilaterally without compensation 	<ul style="list-style-type: none"> Do not force motion Avoid RTC pain with strengthening 	<ul style="list-style-type: none"> Week 16: Functional testing including HHD for IR/ER/Flexion and UE Y-balance Advancement to isotonic exercise per tolerance in all planes, including multiplane exercises provided: <ul style="list-style-type: none"> No compensations during exercise performance Modalities as needed Week 16: FOTO, QuickDASH
Weeks 20+ (Months 5+)	<ul style="list-style-type: none"> Initiate return to sport progression Initiate higher level impact activity General goal for full return to sport at 6 months, depending on progression and sport demands 	<ul style="list-style-type: none"> Focus on form and control during exercise performance Use of appropriate work rest intervals Assess tolerance to activity during, after and at 24 hours after activity 	<ul style="list-style-type: none"> Low level sport specific activity, progressing to higher demand activity Continue with Anaerobic and aerobic interval training Continue with core stability per tolerance <ul style="list-style-type: none"> Multiple planes Stability in all 3 planes of motion Sport specific movements when able Plyometric activities progressing from simple to complex, less load to more load Week 24: FOTO, QuickDASH



OkorohaPA@gmail.com



1801 Inwood Road, 1st Floor
 Dallas, Texas 75390



Office number
 214-645-3300

This protocol is not meant to be prescriptive but a recommendation to guide the rehabilitation process.
 Each patient's progress may vary based on specifics to their injury and procedure.