



Sports Medicine, Joint Preservation & Cartilage Restoration

Kelechi R. Okoroha, MD

HIP, KNEE, SHOULDER, ELBOW

Henry Ford Health System
690 Amsterdam St., Detroit MI
48202 Ph: 313-972-4216
www.KelechiOkorohaMD.com

COMBINED OSTEOCHONDRAL ALLOGRAFT MENISCAL ALLOGRAFT TRANPLANTATION DISTAL FEMORAL OSTEOTOMY

Rehab Protocol

	WEIGHT BEARING	BRACE	ROM	EXERCISES
PHASE I 0-2 weeks	Heel-touch only	Locked in full extension at all times* Off for hygiene and home exercise only	Gentle passive 0-90° CPM 0-90°	Heel slides, quad sets, patellar mobs, SLR, calf pumps at home
PHASE II 2-8 weeks	2-6 weeks: Heel-touch only 6-8 weeks: Advance 25% weekly until full	2-6 weeks: Locked 0-90° Discontinue brace at 6 weeks	Advance as tolerated w/ caution during flexion >90° to protect post horn of meniscus	2-6 weeks: Add side-lying hip and core, advance quad set and stretching** 6-8 weeks: Addition of heel raises, total gym (closed chain), gait normalization, eccentric quads, eccentric hamstrings Advance core, glutes and pelvic stability
PHASE III 8-12 weeks	Full	None	Full	Progress closed chain activities Advance hamstring work, lunges/leg press 0-90° only, proprioception/balance exercises Begin stationary bike
PHASE IV 12-24 weeks	Full	None	Full	Progress Phase III exercises and functional activities: walking lunges, planks, bridges, swiss ball, half-bosu exercises Advance core/glutes and balance
PHASE V 6-9 months	Full	None	Full	Advance all activity w/o impact such as running, jumping, pivoting, sports until cleared by MD

*Brace may be removed for sleeping after first post-operative visit (day 7-14)

**Avoid any tibial rotation for 8 weeks to protect meniscus



OkorohaPA@gmail.com



37000 Woodward Ave., Ste.300, Bloomfield Hills, MI 248-952-9200

14500 Hall Rd., Sterling Hgts. MI 48313 586-247-3714



Secretary
313-651-1926

Other Locations