Henry Ford Health System

690 Amsterdam St., Detroit MI 48202 Ph: 313-972-4216 www.KelechiOkorohaMD.com

MPFL ALLOGRAFT RECONSTRUCTION

Rehab Protocol

	WEIGHT BEARING	BRACE	ROM	EXERCISES
PHASE I 0-2 weeks	As tolerated with brace	On at all times during day and while sleeping** Off for hygiene	0-90°	Heel slides, quad and hamstring sets, patellar mobs, gastroc stretch, SLR in brace, modalities
PHASE II 2-6 weeks	As tolerated with brace	Discontinue at 6 weeks if able to obtain full extension w/o lag*	Maintain full extension and progress flexion	Progress weight bearing flexibility, begin toe raises and closed chain quad work Begin floor-based core and glutes work, balance exercises, hamstring curls and stationary bike
PHASE III 6 weeks - 4 months	Full	None	Full	Advance closed chain quads, progress balance, core/pelvic and stability work Begin elliptical, in-line jogging at 12 weeks under PT supervision
PHASE IV 4-6 months	Full	None	Full	Progress flexibility/strengthening, progression of function: forward/backward running, cutting, grapevine, initiate plyometric program and sport-specific drills @ 16 wks Return to play as tolerated after 16 weeks post-op when cleared by MD

^{*}Brace may be changed to patellar-stabilizing brace (Hely-Weber, Shields, or equivalent) after first post-operative visit



^{**}Brace may be removed for sleeping after first post-operative visit (day 7-10)